

Naive dimensional analysis and truly strong interactions

B. Holdom*

Department of Physics, University of Toronto

Toronto, Ontario, M5S1A7, CANADA

Abstract

We question whether the canonical estimate of a strong coupling, $\alpha \approx 4\pi$, is always appropriate for dynamical symmetry breaking. Our discussion is motivated by the success of naive quark models in describing low energy QCD.

Naive dimensional analysis (NDA) [1] has emerged as a practical tool for estimating the dimensionless parameters appearing in a low energy effective theory when the underlying theory is strongly-coupled. The basic reasonable assumption is that the effective theory should reflect the strongly-interacting nature of the underlying theory, so that loop corrections in the effective theory should be as large as tree level effects. This yields estimates for coefficients of operators in the effective theory. Such estimates turn out to work rather well for low energy QCD, and since the basic idea seems very general, it is natural to apply it to other types of strong interactions.

Central to the application of NDA is an estimate of the size of coupling in a strongly-coupled theory. The canonical estimate is $\alpha \approx 4\pi$, since it is for this coupling that all orders of the loop expansion are expected to be of similar size. We will refer to $\alpha \approx 4\pi$ as a truly strong coupling, and in the absence of any small expansion parameter, the parameters appearing in the low energy effective theory are truly incalculable in perturbation theory. NDA is the best that can be done in the absence of a true nonperturbative calculation.

We will suggest that symmetry breaking physics need not always be truly strong in the following sense. It may be that symmetry breaking physics can at times leave a recognizable imprint on the low energy theory, an imprint not completely obscured

*holdom@utcc.utoronto.ca

by nonperturbative effects and reflecting the details of the underlying theory. We suggest that a coupling which is not truly strong is consistent with the nonperturbative “strongly-coupled” effects underlying the dynamical symmetry breaking. This may be the situation for QCD, such that the success of NDA estimates can be reconciled with the success of naive quark model estimates in low energy QCD. Our discussion does not challenge the basic notion of NDA.

We begin with an example which illustrates the possibility of symmetry breaking physics without a truly strong coupling. We look at a chiral gauge theory and compare NDA with some other standard lore, namely that which follows from the ladder Schwinger-Dyson analysis (LSDA). The latter relies on a one gauge boson approximation to the SD equation. One of the most familiar chiral gauge theories is $SU(5)$ with fermion content $10 + \bar{5}$. At strong enough couplings it is expected that condensates in the channels

$$10 \times 10 \rightarrow \bar{5} \quad \bar{5} \times 10 \rightarrow 5 \quad (1)$$

will occur, since these are the most attractive channels for the respective pairs of fermions [2]. LSDA gives the critical couplings needed for these condensates as $\alpha_c = 5\pi/36 = .44$ and $\alpha_c = 5\pi/27 = .58$ respectively. We may take the first as the critical coupling for the symmetry breaking $SU(5) \rightarrow SU(4)$. (Whether the second condensate is considered to arise from the $SU(5)$ interactions or the unbroken $SU(4)$ interactions is a matter of taste.) Far below the $SU(5)$ breaking scale and the $SU(4)$ confining scale all degrees of freedom have decoupled from the low energy theory except for one fermion. This is the $SU(4)$ singlet component of the original $\bar{5}$ fermion multiplet, the left-handed field ψ_5 , whose masslessness is protected by an unbroken global symmetry [2].

The critical couplings obtained from LSDA are well below $\alpha \approx 4\pi$. LSDA implies that the sum of ladder graphs in specific attractive channels is sufficient to produce dynamical symmetry breaking. The theory is strongly-coupled in the sense that the symmetry breaking effect is nonperturbative; graphs at all orders are contributing. The discrepancy with NDA is simply in the value of the critical coupling in LSDA, which is less than truly strong.

It is tempting to dismiss the naive LSDA results, based as they are on the assumption of a constant coupling. On the other hand we may associate this constant coupling with the value of the true running coupling when smeared or averaged over

the dominant range of momenta in loops. The basic point then is that the LSDA result is presumably reasonable if higher order corrections to the ladder sum are small, which may follow if the coupling is indeed not truly strong. In fact this is consistent with explicit calculations of next order corrections to the LSDA, where relatively small corrections are found [3]. Thus the LSDA results appear to be self-consistent.

Dynamical symmetry breaking in theories which are not truly strong would clearly have implications for the resulting low energy effective theories. In the present example the exchange of a massive $SU(5)$ gauge boson generates the following interaction.

$$\frac{4\pi}{5} \frac{\alpha_c}{M^2} \sum_{n=0}^{\infty} \bar{\psi}_5 \gamma_\mu \psi_5 \left(\frac{\partial^2}{M^2} \right)^n \bar{\psi}_5 \gamma^\mu \psi_5 \quad (2)$$

But by NDA the effective action of the low energy theory should be of the following form, where all dimensionless parameters appearing in the Lagrangian are of order unity [1].

$$S = \int d^4x \frac{\Lambda^4}{(4\pi)^2} \mathcal{L} \left(\frac{4\pi\psi_5}{\Lambda^{\frac{3}{2}}}, \frac{\partial}{\Lambda} \right) \quad (3)$$

The set of terms in (2) could be consistent with (3) only if the following relations were true.

$$M \stackrel{?}{\approx} \Lambda \quad (4)$$

$$\frac{4\pi}{5} \frac{\alpha_c}{M^2} \stackrel{?}{\approx} \frac{(4\pi)^2}{\Lambda^2} \quad (5)$$

If we assume that (4) is true then the NDA estimate of the overall size of the four-fermion operators (RHS of (5)) is 144 times larger than the LSDA estimate (the LHS)! Part of the blame for this large discrepancy surely lies in LSDA, which ignores the fact that the $SU(5)$ coupling is running quite quickly near the $SU(5)$ breaking scale. Thus even though some smeared value of the coupling appropriate for loop integrations may be close to α_c , the α_c appearing in (2) is likely an underestimate. On the other hand for chiral gauge theories with walking couplings, the corresponding effect would be smaller.

The large range in possible estimates for the size of four-fermion operators does not contradict the order-of-magnitude estimates of NDA. But the implication from LSDA, and a coupling which is not truly strong, is that the effect of loop corrections can presumably be quite modest. The specific form of the four-fermion interactions in (2) for example can remain as a clear imprint on the low energy theory.

In the above example the mass of the heavy boson set the scale for the momentum expansion, whereas in the next example involving a heavy fermion the situation is a little different. Consider a standard nonlinear σ -model with a triplet of massless pions coupled to a degenerate fermion doublet of mass m .

$$\begin{aligned}\mathcal{L} &= \frac{f^2}{4} \text{Tr} D_\mu U D^\mu U^\dagger + \bar{\psi}(i\not{D} + \not{V} + A\not{\gamma}_5)\psi - m\bar{\psi}U\psi \\ U(x) &\equiv e^{-\frac{2i\pi(x)\gamma_5}{f}}, \quad V \equiv V_i\sigma_i, \quad \pi \equiv \pi_i\sigma_i \\ D_\mu U &\equiv \partial_\mu U - i(V_\mu + A_\mu)U + iU(V_\mu - A_\mu)\end{aligned}\tag{6}$$

The model has local $SU(2)_L \times SU(2)_R$ symmetry. When used as a model for low energy QCD, color is introduced as a global symmetry where each quark comes in N_c colors. The fermions may be integrated out to yield the low energy theory for the massless pions. We note that this is a theory with two independent mass terms, the ψ mass and the A_μ mass. The pion kinetic term containing the latter cannot be removed due to the infinite renormalization from the fermion loop.

But we are still able to study the finite higher order terms in the momentum expansion of the fermion loop. We consider momentum expansions for the $VV - AA$ two-point function and the electromagnetic pion form factor.

$$\int e^{iqx} \langle \text{TV}_{\mu a}(x) V_{\nu b}(0) - \text{TA}_{\mu a}(x) A_{\nu b}(0) \rangle dx = iF_{V-A}(q^2) f^2 \delta_{ab} (g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2}) \tag{7}$$

$$\langle \pi_a(q_2) | V_b^\mu | \pi_c(q_1) \rangle = i\varepsilon_{abc} F_{V\pi\pi}(q^2) (q_1^\mu + q_2^\mu), \quad q \equiv q_2 - q_1 \tag{8}$$

Both form factors are defined such that $F_{V-A}(0) = F_{V\pi^2}(0) = 1$. The model in fact produces the same result for the two form factors for any q^2 .

$$F_{V-A}(q^2) = F_{V\pi^2}(q^2) = 1 + \frac{N_c}{4\pi^2} \frac{m^2}{f^2} (2 - 2\sqrt{\frac{4m^2 - q^2}{q^2}} \arctan(\sqrt{\frac{q^2}{4m^2 - q^2}})) \tag{9}$$

If we consider the momentum expansion of this result in the form

$$1 + \frac{N_c}{24\pi^2} \frac{q^2}{f^2} \left(1 + \sum_{n=1}^{\infty} \left(a_n \frac{q^2}{m^2} \right)^n \right), \tag{10}$$

we find that the a_n are significantly less than unity, as shown in Fig. (1) for a_n up to high n . Instead of using m as the mass scale in the momentum expansion, a choice $\approx 3m$ would work better for the lower orders, while $2m$ (the threshold for pair production) would work better for the very high orders. We will see that the mass scale appearing in the momentum expansion and its connection to the fermion mass is relevant for the next example, which is closer in spirit to QCD.

The model here will have only one independent mass parameter so that the coupling strength m/f is determined. The independence of f and m in the previous model is related to the infinite renormalization, which in turn is a consequence of the momentum independence of the fermion mass. We will therefore look at a model which incorporates a momentum dependent fermion mass function, which is expected in any case when the fermion mass has a dynamical origin. The minimal model [4] which incorporates the same local chiral gauge symmetries as (6) is obtained by removing the $\text{Tr} D_\mu U D^\mu U^\dagger$ term in (6) and making the following replacement.

$$\begin{aligned} m\bar{\psi}U\psi &\rightarrow \Sigma(x-y)\bar{\psi}(x)\xi(x)X(x,y)\xi(y)\psi(y) \\ \xi(x)^2 &\equiv U(x), \quad X(x,y) \equiv \text{Pe}^{(-i\int_x^y \Gamma_\mu(z)dz^\mu)} \\ \Gamma_\mu &\equiv \frac{i}{2}(\xi(\partial_\mu - iV_\mu - iA_\mu\gamma_5)\xi^\dagger + \xi^\dagger(\partial_\mu - iV_\mu + iA_\mu\gamma_5)\xi) \end{aligned} \quad (11)$$

$\Sigma(x-y)$ is the Fourier transform of $\Sigma(-p^2)$. When the latter function falls for large $-p^2$ in the way appropriate for QCD, the fermion loop generates a finite contribution to the $\text{Tr} D_\mu U D^\mu U^\dagger$ term.

The new quark propagator is $i/(\not{p} - \Sigma(-p^2))$. After expanding the quark loop in external momentum and Wick rotating, we are left with an integral over Euclidean momentum $P^2 = -p^2 > 0$. Since the integral is dominated for $P^2 \approx m^2$ we choose the normalization condition $\Sigma(m^2) = m$ in order for the new propagator to resemble $i/(\not{p} - m)$ in the dominant momentum region. Thus m is again the typical mass parameter of the underlying theory.

The proper normalization of the $\text{Tr} D_\mu U D^\mu U^\dagger$ term yields the Pagels-Stokar [5] relation.

$$f^2 = \frac{N_c}{4\pi^2} \int dP^2 P^2 \frac{\Sigma^2 - P^2 \Sigma \Sigma' / 2}{(P^2 + \Sigma^2)^2} \quad (12)$$

A convenient choice for $\Sigma(P^2)$ is

$$\Sigma(P^2) = \frac{(A+1)m^3}{P^2 + Am^2} \quad (13)$$

and we shall present results for the values $A = 1, 2, 3, 4$.¹ A around 2 or 3 does a good job of reproducing the observed parameters at order p^4 , L_1 – L_{10} , of low energy QCD (after current quark masses are added to the model). Combining (12) and (13) yields the following set of couplings m/f , which do not display extreme sensitivity to

¹Of interest to the modeling of confinement is the fact that the propagator has no pole when $A < 4.83$.

A.

A	1	2	3	4
m/f	4.08	3.77	3.57	3.42

(14)

We again consider the form factors $F_{V\pi^2}(q^2)$ and $F_{V-A}(q^2)$, which are no longer equal. If we expand both form factors in the way suggested by NDA,

$$1 + \sum_{n=1}^{\infty} \left(b_n \frac{N_c q^2}{16\pi^2 f^2} \right)^n \quad (15)$$

we find the following results for the expansion parameters b_n .²

b_n^{V-A}	$n = 1$	2	3	4
$A = 1$	1.42	.92	.81	.80
2	1.16	.78	.66	.63
3	1.05	.73	.61	.56
4	.98	.71	.60	.54

(16)

$b_n^{V\pi^2}$	$n = 1$	2	3
$A = 1$	.90	.70	.69
2	.77	.61	.56
3	.73	.59	.52
4	.70	.58	.52

(17)

We see a tendency for b_n to decrease for increasing n , but to the extent that the b_n are close to unity, the results are fairly consistent with the expectations of NDA for a strongly-interacting, effective theory. Their values would suggest that the fundamental mass scale in the underlying theory is somewhat larger than $4\pi f/\sqrt{N_c}$, and that the underlying theory has coupling somewhat larger than $4\pi/\sqrt{N_c}$. On the other hand, $4\pi f/\sqrt{N_c}$ and $4\pi/\sqrt{N_c}$ are somewhat larger than the mass m and coupling m/f actually appearing in the underlying theory. This is understandable from the previous example, where we learned that the mass of the fermion in the loop is not the appropriate scale for the momentum expansion.

We can now return to the question of how it is that both NDA and naive quark models, such as the one above, are both successful in describing low energy QCD. The point is that what appears to be a strongly-interacting low energy theory is emerging from an underlying theory with a smaller than expected coupling. In fact this coupling, m/f , in the model is very similar to the size of the gauge coupling, $g \approx \sqrt{4\pi}$, which emerges from the ladder Schwinger-Dyson equation analysis of chiral symmetry breaking in QCD. Thus the idea of symmetry breaking arising for couplings

²These numbers are converted from those listed in [6].

less than the truly strong coupling $g \approx 4\pi$ emerges again here, as it did for our first example. Support for this idea appears in the observed form of the chiral Lagrangian of QCD, which seems to display the imprint of some rather trivial physics.

Acknowledgment

I thank M. Luke for a useful discussion. This research was supported in part by the Natural Sciences and Engineering Research Council of Canada.

References

- [1] H. Georgi and A. Manohar, Nucl. Phys. **234B**, 189 (1984); H. Georgi and L. Randall, Nucl. Phys. **B276**, 241 (1986); H. Georgi, Phys. Lett. **B298**, 187 (1993); M. A. Luty, hep-ph/9706235; A. G. Cohen, D. B. Kaplan, A. E. Nelson, hep-ph/9706275.
- [2] S. Raby, S. Dimopoulos, and L. Susskind, Nucl. Phys. **B169**, 373 (1980).
- [3] T. Appelquist., K. Lane, U. Mahanta, Phys. Rev. Lett. **61**, 1553 (1988).
- [4] B. Holdom, Phys. Rev. **D45**, 2534 (1992); J. Terning, Phys. Rev. **D44**, 887 (1991); B. Holdom, J. Terning, and K. Verbeek, Phys. Lett. **B245**, 612 (1990); **B273**, 549E (1991).
- [5] H. Pagels and S. Stokar, Phys. Rev. **D20**, 2947 (1979).
- [6] B. Holdom, Phys. Lett. **B292**, 150 (1992).




Figure (1): The expansion coefficients a_n in (10) asymptotically approaching $1/2^2$.